

# Boolean Coverings of Quantum Observable Structure: A Setting for an Abstract Differential Geometric Mechanism

ELIAS ZAFIRIS

*University of Sofia*

*Faculty of Mathematics and Informatics*

*blvd. James Bourchier, 1164 Sofia*

*Bulgaria*

*e.mail: e.zafiris@fmi.uni-sofia.bg*

## Abstract

We develop the idea of employing localization systems of Boolean coverings, associated with measurement situations, in order to comprehend structures of Quantum Observables. In this manner, Boolean domain observables constitute structure sheaves of coordinatization coefficients in the attempt to probe the Quantum world. Interpretational aspects of the proposed scheme are discussed with respect to a functorial formulation of information exchange, as well as, quantum logical considerations. Finally, the sheaf theoretical construction suggests an operationally intuitive method to develop differential geometric concepts in the quantum regime.

# 1 Introduction

The main guiding idea in our investigation is based on the employment of objects belonging to the Boolean species of observable structure, as covers, for the understanding of the objects belonging to the quantum species of observable structure. The language of Category theory [1, 2] proves to be suitable for the implementation of this idea in a universal way. The conceptual essence of this scheme is the development of a sheaf theoretical perspective [3, 4] on Quantum observable structures.

The physical interpretation of the categorical framework makes use of the analogy with geometric manifold theory. Namely, it is associated with the development of a Boolean manifold picture, that takes place through the identification of Boolean charts in systems of localization for quantum event algebras with reference frames, relative to which the results of measurements can be coordinatized. In this sense, any Boolean chart in a localization system covering a quantum algebra of events, corresponds to a set of Boolean events which become realizable in the experimental context of a measurement situation. This identification amounts to the introduction of a relativity principle in Quantum theory, suggesting a contextual interpretation of its descriptive apparatus.

In quantum logical approaches the notion of event, associated with the measurement of an observable, is taken to be equivalent to a proposition describing the behavior of a physical system. This formulation of Quantum theory is based on the identification of propositions with projection operators on a complex Hilbert space. In this sense, the Hilbert space formalism of Quantum theory associates events with closed subspaces of a separable, complex Hilbert space corresponding to a quantum system. Then, the quantum event algebra is identified with the lattice of closed subspaces of the Hilbert space, ordered by inclusion and carrying an orthocomplementation operation which is given by the orthogonal complements of the closed subspaces [5-6]. Equivalently it is isomorphic to the partial Boolean algebra of closed subspaces of the Hilbert space of the system, or alternatively the partial Boolean algebra of projection operators of the system [7].

We argue that the set theoretical axiomatizations of quantum observable structures hides the intrinsic significance of Boolean localizing systems in the formation of these structures. Moreover, the operational procedures followed in quantum measurement are based explicitly in the employment of appropriate Boolean environments. The construction of these contexts of observation are related with certain abstractions and can be metaphorically considered as pattern recognition arrangements. In the categorical language we adopt, we can explicitly associate them with appropriate Boolean coverings of the

structure of quantum events. In this way, the real significance of a quantum structure proves to be, not at the level of events, but at the level of gluing together observational contexts.

The main thesis of this paper, is that, the objectification of a quantum observable structure takes place through Boolean reference frames, that can be pasted together using category theoretical means. Contextual topos theoretical approaches to quantum structures have been considered, from a different viewpoint in [8,9], and discussed in [10-12].

In Section 2, we define event and observable structures in a category theoretical language. In Section 3, we introduce the functorial concepts of Boolean coordinatizations and Boolean observable presheaves, and also, develop the idea of fibrations over Boolean observables. In Section 4, we prove the existence of an adjunction between the topos of presheaves of Boolean observables and the category of Quantum observables. In Section 5, we define systems of localization for measurement of observables over a quantum event and analyze their physical significance. In Section 6 we talk about isomorphic representations of quantum algebras in terms of Boolean localization systems using the adjunction established. In Section 7, we examine the consequences of the scheme related to the interpretation of the logic of quantum propositions. In Section 8, we discuss the implications of covering systems, in relation to, the possibility of development of a differential geometric machinery, suitable for the quantum regime. Finally, we summarize the conclusions in Section 9.

## 2 Event and Observable Structures as Categories

A **Classical event structure** is a category, denoted by  $\mathcal{B}$ , which is called the category of Boolean event algebras. Its objects are Boolean algebras of events, and its arrows are Boolean algebraic homomorphisms.

A **Quantum event structure** is a category, denoted by  $\mathcal{L}$ , which is called the category of Quantum event algebras.

Its objects are Quantum algebras of events, that is, partially ordered sets of Quantum events, endowed with a maximal element 1, and with an operation of orthocomplementation  $[-]^* : L \longrightarrow L$ , which satisfy, for all  $l \in L$  the following conditions: [a]  $l \leq 1$ , [b]  $l^{**} = l$ , [c]  $l \vee l^* = 1$ , [d]  $l \leq \acute{l} \Rightarrow \acute{l}^* \leq l^*$ , [e]  $l \perp \acute{l} \Rightarrow l \vee \acute{l} \in L$ , [g]  $l \vee \acute{l} = 1, l \wedge \acute{l} = 0 \Rightarrow l = \acute{l}^*$ , where  $0 := 1^*$ ,  $l \perp \acute{l} := l \leq \acute{l}^*$ , and the operations of meet  $\wedge$  and join  $\vee$  are defined as usually.

Its arrows are Quantum algebraic homomorphisms, that is maps  $L \xrightarrow{H} K$ ,

which satisfy, for all  $k \in K$  the following conditions: [a]  $H(1) = 1$ , [b]  $H(k^*) = [H(k)]^*$ , [c]  $k \leq \acute{k} \Rightarrow H(k) \leq H(\acute{k})$ , [d]  $k \perp \acute{k} \Rightarrow H(k \vee \acute{k}) \leq H(k) \vee H(\acute{k})$ .

The notion of observable corresponds to a physical quantity that can be measured in the context of an experimental arrangement. In any measurement situation, the propositions that can be made concerning a physical quantity are of the following type: the value of the physical quantity lies in some Borel set of the real numbers. The proposition that the value of a physical quantity lies in a Borel set of the real line corresponds to an event as it is apprehended by an observer using his measuring instrument. In this sense, we construct a mapping from the domain of Borel sets of the real line to the event structure, that captures precisely the notion of observable. In suggestive picture the real line, endowed with its Borel structure, plays the role of a shaping object, that schematizes the event algebra of an observed system, by projecting into it its structure.

Observables can be formulated in the environment of categories as follows:

A **Quantum observable space structure** is a category, denoted by  $\mathcal{OB}$ , which is called the category of spaces of Quantum observables.

Its objects are the sets  $\Omega$  of real-valued observables on a quantum event algebra  $L$ , where each observable  $\Xi$  is defined to be an algebraic homomorphism from the Borel algebra of the real line  $Bor(\mathbf{R})$ , to the quantum event algebra  $L$ .

$$\Xi : Bor(\mathbf{R}) \rightarrow L$$

such that the following conditions are satisfied:

[i]  $\Xi(\emptyset) = 0, \Xi(\mathbf{R}) = 1$ , [ii]  $E \cap F = \emptyset \Rightarrow \Xi(E) \perp \Xi(F)$ , for  $E, F \in Bor(\mathbf{R})$ , [iii]  $\Xi(\bigcup_n E_n) = \bigvee_n \Xi(E_n)$ , where  $E_1, E_2, \dots$  sequence of mutually disjoint Borel sets of the real line. Its arrows are the quantum observable spaces homomorphisms.

If  $L$  is isomorphic with the orthocomplemented lattice of orthogonal projections on a Hilbert space, then it follows from von Neumann's spectral theorem that the observables are in 1-1 correspondence with the hypermaximal Hermitian operators on the Hilbert space.

Using the information encoded in the categories of Quantum event algebras  $\mathcal{L}$ , and spaces of Quantum observables  $\mathcal{OB}$ , it is possible to construct a new category, called the category of Quantum observables, defined as follows:

A **Quantum observable structure** is a category, denoted by  $\mathcal{O}_Q$ , which is called the category of Quantum observables.

Its objects are the quantum observables  $\Xi : Bor(\mathbf{R}) \rightarrow L$  and its arrows  $\Xi \longrightarrow \Theta$  are the commutative triangles [Diagram 1], or equivalently the quantum algebraic homomorphisms  $L \xrightarrow{H} K$  in  $\mathcal{L}$ , such that  $\Theta = H \circ \Xi$  in




Diagram 1




Diagram 2

[Diagram 1] is again a quantum observable.

Correspondingly, a **Boolean observable structure** is a category, denoted by  $\mathcal{O}_B$ , which is called the category of Boolean observables.

Its objects are the Boolean observables  $\xi : \text{Bor}(\mathbf{R}) \rightarrow B$  and its arrows are the Boolean algebraic homomorphisms  $B \xrightarrow{h} C$  in  $\mathcal{B}$ , such that  $\theta = h \circ \xi$  in [Diagram 2] is again a Boolean observable.

### 3 Functorial Formulation of Observable Structures

#### 3.1 Presheaves of Boolean Observables

If  $\mathcal{O}_B^{op}$  is the opposite category of  $\mathcal{O}_B$ , then  $\mathbf{Sets}^{\mathcal{O}_B^{op}}$  denotes the functor category of presheaves on Boolean observables. Its objects are all functors  $\mathbf{X} : \mathcal{O}_B^{op} \rightarrow \mathbf{Sets}$ , and its morphisms are all natural transformations between such functors. Each object  $\mathbf{X}$  in this category is a contravariant set-valued functor on  $\mathcal{O}_B$ , called a presheaf on  $\mathcal{O}_B$ .

For each Boolean observable  $\xi$  of  $\mathcal{O}_B$ ,  $\mathbf{X}(\xi)$  is a set, and for each arrow  $f : \theta \rightarrow \xi$ ,  $\mathbf{X}(f) : \mathbf{X}(\xi) \rightarrow \mathbf{X}(\theta)$  is a set function. If  $\mathbf{X}$  is a presheaf on  $\mathcal{O}_B$  and  $x \in \mathbf{X}(O)$ , the value  $\mathbf{X}(f)(x)$  for an arrow  $f : \theta \rightarrow \xi$  in  $\mathcal{O}_B$  is called the restriction of  $x$  along  $f$  and is denoted by  $\mathbf{X}(f)(x) = x \circ f$ .

Each object  $\xi$  of  $\mathcal{O}_B$  gives rise to a contravariant Hom-functor  $y[\xi] := \text{Hom}_{\mathcal{O}_B}(-, \xi)$ . This functor defines a presheaf on  $\mathcal{O}_B$ . Its action on an object  $\theta$  of  $\mathcal{O}_B$  is given by

$$y[\xi](\theta) := \text{Hom}_{\mathcal{O}_B}(\theta, \xi)$$

whereas its action on a morphism  $\eta \xrightarrow{w} \theta$ , for  $v : \theta \longrightarrow \xi$  is given by

$$y[\xi](w) : \text{Hom}_{\mathcal{O}_B}(\theta, \xi) \longrightarrow \text{Hom}_{\mathcal{O}_B}(\eta, \xi)$$

$$y[\xi](w)(v) = v \circ w$$

Furthermore  $y$  can be made into a functor from  $\mathcal{O}_B$  to the contravariant functors on  $\mathcal{O}_B$

$$y : \mathcal{O}_B \longrightarrow \mathbf{Sets}^{\mathcal{O}_B^{op}}$$

such that  $\xi \mapsto \text{Hom}_{\mathcal{O}_B}(-, \xi)$ . This is an embedding and it is a full and faithful functor.

The functor category of presheaves on Boolean observables  $\mathbf{Sets}^{\mathcal{O}_B^{op}}$ , provides an instantiation of a structure known as topos. A topos exemplifies a well defined notion of variable set. It can be conceived as a local mathematical framework corresponding to a generalized model of set theory or as a generalized space. Moreover it provides a natural example of a many-valued truth structure, which remarkably is not ad hoc, but reflects genuine constraints of the surrounding universe. The study of the truth value structure associated with the topos of presheaves of Boolean observables and its significance for a quantum logical interpretation of the proposed categorical scheme will be the subject of a separate paper. Some ideas related to this direction are discussed in Section 9.

### 3.2 The Grothendieck Fibration Technique

Since  $\mathcal{O}_B$  is a small category, there is a set consisting of all the elements of all the sets  $\mathbf{X}(\xi)$ , and similarly there is a set consisting of all the functions  $\mathbf{X}(f)$ . This observation regarding  $\mathbf{X} : \mathcal{O}_B^{op} \longrightarrow \mathbf{Sets}$  permits us to take the disjoint union of all the sets of the form  $\mathbf{X}(\xi)$  for all objects  $\xi$  of  $\mathcal{O}_B$ . The elements of this disjoint union can be represented as pairs  $(\xi, x)$  for all objects  $\xi$  of  $\mathcal{O}_B$  and elements  $x \in \mathbf{X}(\xi)$ . Thus the disjoint union of sets is made by labeling the elements. Now we can construct a category whose set of objects is the disjoint union just mentioned. This structure is called the category of elements of the presheaf  $\mathbf{X}$ , denoted by  $\mathbf{G}(\mathbf{X}, \mathcal{O}_B)$ . Its objects are all pairs  $(\xi, x)$ , and its morphisms  $(\acute{\xi}, \acute{x}) \longrightarrow (\xi, x)$  are those morphisms  $u : \acute{\xi} \longrightarrow \xi$  of  $\mathcal{O}_B$  for which  $xu = \acute{x}$ . Projection on the second coordinate

$$\begin{array}{ccc}
& \mathbf{G}(\mathbf{X}, \mathcal{O}_B) & \\
& \downarrow \mathbf{G}_\mathbf{X} & \\
\mathcal{O}_B & \xrightarrow{\mathbf{X}} & \mathbf{Sets}
\end{array}$$

Diagram 3

$$\begin{array}{ccc}
& \text{Bor}(R) & \\
& \swarrow \xi \quad \searrow \Xi & \\
\mathbf{A}(B_\Xi) & \xrightarrow{[\psi_B]_\Xi} & L
\end{array}$$

Diagram 4

of  $\mathbf{G}(\mathbf{X}, \mathcal{O}_B)$ , defines a functor  $\mathbf{G}_\mathbf{X} : \mathbf{G}(\mathbf{X}, \mathcal{O}_B) \longrightarrow \mathcal{O}_B$ .  $\mathbf{G}(\mathbf{X}, \mathcal{O}_B)$  together with the projection functor  $\mathbf{G}_\mathbf{X}$  is called the split discrete fibration induced by  $\mathbf{X}$ , and  $\mathcal{O}_B$  is the base category of the fibration. We note that the fibration is discrete because the fibers are categories in which the only arrows are identity arrows. If  $\xi$  is a Boolean observable object of  $\mathcal{O}_B$ , the inverse image under  $\mathbf{G}_\mathbf{P}$  of  $\xi$  is simply the set  $\mathbf{X}(\xi)$ , although its elements are written as pairs so as to form a disjoint union. The instantiation of the fibration induced by  $\mathbf{P}$ , is an application of the general Grothendieck construction [13].

### 3.3 Boolean Modeling Functor

We define a modeling or coordinatisation functor,  $\mathbf{A} : \mathcal{O}_B \longrightarrow \mathcal{O}_Q$ , which assigns to Boolean observables in  $\mathcal{O}_B$  (that plays the role of the model category) the underlying Quantum observables from  $\mathcal{O}_Q$ , and to Boolean homomorphisms the underlying quantum algebraic homomorphisms. Hence  $\mathbf{A}$  acts as a forgetful functor, forgetting the extra Boolean structure of  $\mathcal{O}_B$ .

Equivalently the coordinatisation functor can be characterized as,  $\mathbf{A} : \mathcal{B} \longrightarrow \mathcal{L}$ , which assigns to Boolean event algebras in  $\mathcal{B}$  the underlying quantum event algebras from  $\mathcal{L}$ , and to Boolean homomorphisms the underlying quantum algebraic homomorphisms, such that [Diagram 4] commutes:

$$\begin{array}{ccc}
\mathbf{X}(\xi) & \xrightarrow{\tau_\xi} & \text{Hom}_{\mathcal{O}_Q}(\mathbf{A}(\xi), \Xi) \\
\mathbf{X}(u) \downarrow & & \downarrow \mathbf{A}(u) \\
\mathbf{X}(\xi') & \xrightarrow{\tau_{\xi'}} & \text{Hom}_{\mathcal{O}_Q}(\mathbf{A}(\xi'), \Xi)
\end{array}$$

Diagram 5

### 3.4 Functorial Relation of Event with Observable Algebras

The categories of Event algebras and Observables are related functorially as follows: Under the action of a modeling functor,  $\text{Bor}\mathbf{R}$  may be considered as an object of  $\mathcal{L}$ . Hence, it is possible to construct the covariant representable functor  $\mathbf{F} : \mathcal{L} \rightarrow \mathbf{Sets}$ , defined by  $\mathbf{F} = \text{Hom}_{\mathcal{L}}(\text{Bor}\mathbf{R}, -)$ . The application of the fibration technique on the functor  $\mathbf{F}$  provides the category of elements of this functor, which is the category of all arrows in  $\mathcal{L}$  from the object  $\text{Bor}\mathbf{R}$ , characterized equivalently as the comma category  $[\text{Bor}\mathbf{R}/\mathcal{L}]$ . We conclude that the category of Quantum observables  $\mathcal{O}_Q$  is actually the comma category  $[\text{Bor}\mathbf{R}/\mathcal{L}]$  or equivalently the category of elements of the functor  $\mathbf{F} = \text{Hom}_{\mathcal{L}}(\text{Bor}\mathbf{R}, -)$ . Analogous comments hold for the category of Boolean observables.

## 4 Adjointness between Presheaves of Boolean Observables and Quantum Observables

We consider the category of quantum observables  $\mathcal{O}_Q$  and the modeling functor  $\mathbf{A}$ , and we define the functor  $\mathbf{R}$  from  $\mathcal{O}_Q$  to the topos of presheaves given by

$$\mathbf{R}(\Xi) : \xi \mapsto \text{Hom}_{\mathcal{O}_Q}(\mathbf{A}(\xi), \Xi)$$

A natural transformation  $\tau$  between the topos of presheaves on the category of Boolean observables  $\mathbf{X}$  and  $\mathbf{R}(\Xi)$ ,  $\tau : \mathbf{X} \rightarrow \mathbf{R}(\Xi)$  is a family  $\tau_\xi$  indexed by Boolean observables  $\xi$  of  $\mathcal{O}_B$  for which each  $\tau_\xi$  is a map

$$\tau_\xi : \mathbf{X}(\xi) \rightarrow \text{Hom}_{\mathcal{O}_Q}(\mathbf{A}(\xi), \Xi)$$

of sets, such that the diagram of sets [Diagram 5] commutes for each Boolean homomorphism  $u : \xi' \rightarrow \xi$  of  $\mathcal{O}_B$ .



$$\begin{array}{ccc}
\mathbf{A}(\xi) \equiv \mathbf{A} \circ \mathbf{G}_{\mathbf{X}}(\xi, p) & & \\
\uparrow \mathbf{A}(u) & \nearrow \tau_{\xi}(p) & \\
& & \Xi \\
& \nwarrow \hat{\tau}_{\xi}(\hat{p}) & \\
\mathbf{A}(\xi') \equiv \mathbf{A} \circ \mathbf{G}_{\mathbf{X}}(\xi', \hat{p}) & \uparrow u_* &
\end{array}$$

Diagram 6

If we make use of the category of elements of the Boolean observables-variable set  $X$ , being an object in the topos of presheaves, then the map  $\tau_{\xi}$ , defined above, can be characterized as:

$$\tau_{\xi} : (\xi, p) \rightarrow \text{Hom}_{\mathcal{O}_Q}(\mathbf{A} \circ G_{\mathbf{X}}(\xi, p), \Xi)$$

Equivalently such a  $\tau$  can be seen as a family of arrows of  $\mathcal{O}_Q$  which is being indexed by objects  $(\xi, p)$  of the category of elements of the presheaf of Boolean observables  $\mathbf{X}$ , namely

$$\{\tau_{\xi}(p) : \mathbf{A}(\xi) \rightarrow \Xi\}_{(\xi, p)}$$

From the perspective of the category of elements of  $\mathbf{X}$ , the condition of the commutativity of [Diagram 5] is equivalent with the condition that for each Boolean homomorphism  $u : \xi' \rightarrow \xi$  of  $\mathcal{O}_B$ , [Diagram 6] commutes.

From [Diagram 6] we can see that the arrows  $\tau_{\xi}(p)$  form a cocone from the functor  $\mathbf{A} \circ G_{\mathbf{X}}$  to the quantum observable algebra object  $\Xi$ . Making use of the definition of the colimit, we conclude that each such cocone emerges by the composition of the colimiting cocone with a unique arrow from the colimit  $\mathbf{LX}$  to the quantum observable object  $\Xi$ . In other words, there is a bijection which is natural in  $\mathbf{X}$  and  $\Xi$

$$\text{Nat}(\mathbf{X}, \mathbf{R}(\Xi)) \cong \text{Hom}_{\mathcal{O}_Q}(\mathbf{LX}, \Xi)$$

From the above bijection we are driven to the conclusion that the functor  $\mathbf{R}$  from  $\mathcal{O}_Q$  to the topos of presheaves given by

$$\mathbf{R}(\Xi) : \xi \mapsto \text{Hom}_{\mathcal{O}_Q}(\mathbf{A}(\xi), \Xi)$$

$$\begin{array}{ccc}
Nat(\mathbf{X}, \mathbf{R}(\Xi)) & \xrightarrow{\mathbf{r}} & Hom_{\mathcal{O}_Q}(\mathbf{LX}, \Xi) \\
\parallel & & \parallel \\
Nat(\mathbf{X}, \mathbf{R}(\Xi)) & \xleftarrow{\mathbf{l}} & Hom_{\mathcal{O}_Q}(\mathbf{LX}, \Xi)
\end{array}$$

Diagram 7

has a left adjoint  $\mathbf{L} : \mathbf{Sets}^{\mathcal{O}_B^{op}} \rightarrow \mathcal{O}_Q$ , which is defined for each presheaf of Boolean observables  $\mathbf{X}$  in  $\mathbf{Sets}^{\mathcal{O}_B^{op}}$  as the colimit

$$\mathbf{L}(\mathbf{X}) = Colim\{\mathbf{G}(\mathbf{X}, \mathcal{O}_B) \xrightarrow{\mathbf{G}_X} \mathcal{O}_B \xrightarrow{\mathbf{A}} \mathcal{O}_Q\}$$

Consequently there is a **pair of adjoint functors**  $\mathbf{L} \dashv \mathbf{R}$  as follows:

$$\mathbf{L} : \mathbf{Sets}^{[[Bor\mathbf{R}/B]]^{op}} \xrightleftharpoons{\quad} [Bor\mathbf{R}/\mathcal{L}] : \mathbf{R}$$

The adjunction, which will be the main interpretational tool in the proposed scheme, consists of the functors  $\mathbf{L}$  and  $\mathbf{R}$ , called left and right adjoints with respect to each other respectively, [Diagram 7], as well as, the natural bijection

$$Nat(\mathbf{P}, \mathbf{R}(\Xi)) \cong Hom_{[Bor\mathbf{R}/\mathcal{L}]}(\mathbf{LX}, \Xi)$$

As an application we may use as  $\mathbf{X}$  the representable presheaf of the topos of Boolean observables  $\mathbf{y}[\xi]$ . Then, the bijection defining the adjunction takes the form:

$$Nat(\mathbf{y}[\xi], \mathbf{R}(\Xi)) \cong Hom_{\mathcal{O}_Q}(\mathbf{Ly}[\xi], \Xi)$$

Because the functor  $\mathbf{X} = \mathbf{y}[\xi]$  is representable, the corresponding category of elements  $\mathbf{G}(\mathbf{y}[\xi], \mathcal{O}_B)$  has a terminal object, that is, the element  $1 : \xi \longrightarrow \xi$  of  $\mathbf{y}[\xi](\xi)$ . Therefore, the colimit of the composite  $\mathbf{A} \circ \mathbf{G}_{\mathbf{y}[\xi]}$  is going to be just the value of  $\mathbf{A} \circ \mathbf{G}_{\mathbf{y}[\xi]}$  on the terminal object. Thus, we have

$$\mathbf{Ly}[\xi](\xi) \cong \mathbf{A} \circ \mathbf{G}_{\mathbf{y}[\xi]}(\xi, 1_\xi) = \mathbf{A}(\xi)$$

In this way we provide a characterization of  $\mathbf{A}(\xi)$  as the colimit of the representable presheaf on the category of Boolean observables.

Furthermore, the categorical syntax provides a representation of a colimit as a coequalizer of a coproduct. This representation is significant for the purposes of the present scheme, because it reveals the fact that the left adjoint functor of the adjunction is like the tensor product  $-\otimes_{[Bor\mathbf{R}/B]}\mathbf{A}$ .

$$\begin{array}{ccc}
& \mathcal{O}_B & \\
& \downarrow \text{y} & \searrow \mathbf{A} \\
& \text{Sets}^{\mathcal{O}_B^{op}} \cdots \mathbf{L} \cdots & \mathcal{O}_Q
\end{array}$$

Diagram 8

$$\coprod_{v:\xi \rightarrow \xi} \mathbf{A}(\xi) \xrightarrow[\eta]{\zeta} \coprod_{(\xi,p)} \mathbf{A}(\xi) \xrightarrow{\chi} \mathbf{X} \otimes_{\mathcal{O}_B} \mathbf{A}$$

Diagram 9

More specifically, the coequalizer representation of the colimit  $\mathbf{LX}$  has the form of [Diagram 9], where, the second coproduct is over all the objects  $(\xi, p)$  with  $p \in \mathbf{X}(\xi)$  of the category of elements, while the first coproduct is over all the maps  $v : (\xi', p') \longrightarrow (\xi, p)$  of that category, so that  $v : \xi' \longrightarrow \xi$  and the condition  $pv = p'$  is satisfied. The elements of  $\mathbf{X} \otimes_{\mathcal{O}_B} \mathbf{A}$ , considered as a set endowed with a quantum algebraic structure, are all of the form  $\chi(p, q)$ , or in a suggestive notation,

$$\chi(p, q) = p \otimes q, \quad p \in \mathbf{X}(\xi), q \in \mathbf{A}(\xi)$$

satisfying, the coequalizer condition  $pv \otimes q = p \otimes vq$ .

## 5 System Of Measurement Localizations For Quantum Observables

The conceptual basis underlying the notion of a system of localizations for a quantum observable, which will be defined subsequently, is an implication of the categorical principle according to which, the quantum object  $\Xi$  in  $\mathcal{O}_Q$  is possible to be comprehended by means of appropriate covering maps  $\xi \longrightarrow \Xi$  having as their domains, locally defined Boolean observables  $\xi$  in  $\mathcal{O}_B$ . It is obvious that any single map from any modeling Boolean observable to a quantum observable is not sufficient to determine it entirely, and hence, it is a priori destined to contain only a limited amount of information about it. This problem may be tackled, only if, we employ many structure preserving maps from the modeling Boolean observables to a quantum observable simultaneously, so as to cover it completely. In turn, the information available about each map of the specified covering may be used to determine

the quantum observable itself. In this case, we say that, the family of such maps generate a system of prelocalizations for a quantum observable. We can formalize these intuitive ideas as follows:

A **system of prelocalizations** for quantum observable  $\Xi$  in  $\mathcal{O}_Q$  is a subfunctor of the Hom-functor  $\mathbf{R}(\Xi)$  of the form  $\mathbf{S} : \mathcal{O}_B^{op} \rightarrow \mathbf{Sets}$ , namely for all  $\xi$  in  $\mathcal{O}_B$  it satisfies  $\mathbf{S}(\xi) \subseteq [\mathbf{R}(\Xi)](\xi)$ . Hence a system of prelocalizations for quantum observable  $\Xi$  in  $\mathcal{O}_Q$  is a set  $\mathbf{S}(\xi)$  of quantum algebraic homomorphisms of the form

$$\psi_\xi : \mathbf{A}(\xi) \longrightarrow \Xi, \quad \xi \in \mathcal{O}_B$$

such that  $\langle \psi_\xi : \mathbf{A}(\xi) \longrightarrow \Xi \text{ in } \mathbf{S}(\xi), \text{ and } \mathbf{A}(v) : \mathbf{A}(\xi') \rightarrow \mathbf{A}(\xi) \text{ in } \mathcal{O}_Q \text{ for } v : \xi' \rightarrow \xi \text{ in } \mathcal{O}_B, \text{ implies } \psi_\xi \circ \mathbf{A}(v) : \mathbf{A}(\xi') \longrightarrow \Xi \text{ in } \mathbf{S}(\xi') \rangle$ .

According to the above definition, the functional role of the Hom-functor  $\mathbf{R}(\Xi)$  is equivalent to depicting a set of algebraic homomorphisms, in order to provide local coverings of a quantum observable by coordinatizing Boolean objects. We may characterize the maps  $\psi_\xi : \mathbf{A}(\xi) \longrightarrow \Xi, \quad \xi \in \mathcal{O}_B$  in a system of prelocalizations for quantum observable  $\Xi$  as Boolean domain covers. Their domains  $B_\Xi$  provide Boolean coefficients, associated with measurement situations. The introduction of the notion of a system of prelocalizations is forced on the basis of operational physical arguments. According to Kochen-Specker theorem it not possible to understand completely a quantum mechanical system with the use of a single system of Boolean devices. On the other side, in every concrete experimental context, the set of events that have been actualized in this context forms a Boolean algebra. Consequently, any Boolean domain object  $(B_\Xi, [\psi_B]_\Xi : \mathbf{A}(B_\Xi) \longrightarrow L)$  in a system of prelocalizations for quantum event algebra, making [Diagram 10] commutative, corresponds to a set of Boolean events that become actualized in the experimental context of B. These Boolean objects play the role of localizing devices in a quantum event structure, that are induced by measurement situations. The above observation is equivalent to the statement that a measurement-induced Boolean algebra serves as a reference frame, in a topos-theoretical environment, relative to which a measurement result is being coordinatized.

A family of Boolean observable covers  $\psi_\xi : \mathbf{A}(\xi) \longrightarrow \Xi, \quad \xi \in \mathcal{O}_B$  is the generator of the system of prelocalization  $\mathbf{S}$  iff this system is the smallest among all that contains that family. It is evident that a quantum observable, and correspondingly the quantum event algebra over which it is defined, can have many systems of measurement prelocalizations, that, remarkably, form an ordered structure. More specifically, systems of prelocalization constitute a partially ordered set under inclusion. We note that the minimal system is the empty one, namely  $\mathbf{S}(\xi) = \emptyset$  for all  $\xi \in \mathcal{O}_B$ , whereas the maximal

$$\begin{array}{ccc}
\text{Bor}(\mathbf{R}) & \xrightarrow{\xi} & \mathbf{A}(\dot{B}_\Xi) \\
\downarrow \xi & \searrow \Xi & \downarrow [\psi_{\dot{B}}]_\Xi \\
\mathbf{A}(B_\Xi) & \xrightarrow{[\psi_B]_\Xi} & L
\end{array}$$

Diagram 10

$$\begin{array}{ccccc}
\mathbf{T} & & & & \\
& \searrow u & & \searrow h & \\
& & \mathbf{A}(\xi) \times_{\Xi} \mathbf{A}(\xi') & \xrightarrow{\psi_{\xi, \xi'}} & \mathbf{A}(\xi) \\
& \searrow g & \downarrow \psi_{\xi, \xi} & & \downarrow \psi_{\xi} \\
& & \mathbf{A}(\xi') & \xrightarrow{\psi_{\xi'}} & \Xi
\end{array}$$

Diagram 11

system is the Hom-functor  $\mathbf{R}(\Xi)$  itself, or equivalently, all quantum algebraic homomorphisms  $\psi_{\xi} : \mathbf{A}(\xi) \longrightarrow \Xi$ .

The transition from a system of prelocalizations to a system of localizations for a quantum observable, can be realized if certain compatibility conditions are satisfied on the overlap of the modeling Boolean domain covers. In order to accomplish this it is necessary to introduce the categorical concept of pullback in  $\mathcal{O}_Q$  [Diagram 15].

The pullback of the Boolean domain covers  $\psi_{\xi} : \mathbf{A}(\xi) \longrightarrow \Xi, \xi \in \mathcal{O}_B$  and  $\psi_{\xi'} : \mathbf{A}(\xi') \longrightarrow \Xi, \xi' \in \mathcal{O}_B$  with common codomain the quantum observable  $\Xi$ , consists of the object  $\mathbf{A}(\xi) \times_{\Xi} \mathbf{A}(\xi')$  and two arrows  $\psi_{\xi, \xi'}$  and  $\psi_{\xi', \xi}$ , called projections, as shown in [Diagram 11]. The square commutes and for any object  $T$  and arrows  $h$  and  $g$  that make the outer square commute, there is a unique  $u : T \longrightarrow \mathbf{A}(\xi) \times_{\Xi} \mathbf{A}(\xi')$  that makes the whole diagram commutative. Hence we obtain the condition:

$$\psi_{\xi'} \circ g = \psi_{\xi} \circ h$$

We emphasize that if  $\psi_\xi$  and  $\psi_{\xi'}$  are injective maps, then their pullback is isomorphic with the intersection  $\mathbf{A}(\xi) \cap \mathbf{A}(\xi')$ . Then we can define the pasting map, which is an isomorphism, as follows:

$$\Omega_{\xi,\xi'} : \psi_{\xi\xi'}(\mathbf{A}(\xi) \times_{\Xi} \mathbf{A}(\xi')) \longrightarrow \psi_{\xi\xi'}(\mathbf{A}(\xi) \times_{\Xi} \mathbf{A}(\xi'))$$

by putting

$$\Omega_{\xi,\xi} = \psi_{\xi\xi} \circ \psi_{\xi\xi}^{-1}$$

Then we have the following conditions:

$$\begin{aligned} \Omega_{\xi,\xi} &= 1_\xi & 1_\xi &:= id_\xi \\ \Omega_{\xi,\xi} \circ \Omega_{\xi,\xi'} &= \Omega_{\xi,\xi'} & \text{if } \mathbf{A}(\xi) \cap \mathbf{A}(\xi') \cap \mathbf{A}(\xi') &\neq 0 \\ \Omega_{\xi,\xi} &= \Omega_{\xi,\xi} & \text{if } \mathbf{A}(\xi) \cap \mathbf{A}(\xi) &\neq 0 \end{aligned}$$

The pasting map assures that  $\psi_{\xi\xi}(\mathbf{A}(\xi) \times_{\Xi} \mathbf{A}(\xi'))$  and  $\psi_{\xi\xi'}(\mathbf{A}(\xi) \times_{\Xi} \mathbf{A}(\xi'))$  are going to cover the same part of the quantum observable in a compatible way.

Given a system of measurement prelocalizations for quantum observable  $\Xi \in \mathcal{O}_Q$ , and correspondingly for the Quantum event algebra over which it is defined, we call it a **system of localizations**, iff the above compatibility conditions are satisfied, and moreover, the quantum algebraic structure is preserved.

We assert that the above compatibility conditions provide the necessary relations for understanding a system of measurement localizations for a quantum observable as a structure sheaf or sheaf of Boolean coefficients consisting of local Boolean observables. This is connected to the fact that systems of measurement localizations are actually subfunctors of the representable Hom-functor  $\mathbf{R}(\Xi)$  of the form  $\mathbf{S} : \mathcal{O}_B^{op} \rightarrow \mathbf{Sets}$ , namely for all  $\xi$  in  $\mathcal{O}_B$  satisfy  $\mathbf{S}(\xi) \subseteq [\mathbf{R}(\Xi)](\xi)$ . In this sense the pullback compatibility conditions express gluing relations on overlaps of Boolean domain covers and convert a presheaf subfunctor of the Hom-functor into a sheaf.

The concept of sheaf expresses exactly the pasting conditions that local modeling objects have to satisfy, namely, the way by which local data, providing Boolean coefficients obtained in measurement situations, can be collated. We stress the point that the transition from locally defined properties to global consequences happens via a compatible family of elements over a covering system of the complex object. A covering, or equivalently a localization system of the global, complex object, being a Quantum observable structure in the present scheme, can be viewed as providing a decomposition of that object into simpler modeling objects.

$$\begin{array}{ccc}
\Pi_{v:\xi \rightarrow \xi} \mathbf{A}(\xi) & \xrightarrow[\eta]{\zeta} & \Pi_{(\xi,p)} \mathbf{A}(\xi) \rightarrow [\mathbf{R}(\Xi)](-) \otimes_{\mathcal{O}_B} \mathbf{A} \\
& & \searrow \quad \downarrow \epsilon_{\Xi} \\
& & \Xi
\end{array}$$

Diagram 12

The comprehension of a measurement localization system as a sheaf of Boolean coefficients permits the conception of a Quantum observable (or of its associated quantum event algebra) as a generalized manifold, obtained by pasting the  $\psi_{\xi\xi}(\mathbf{A}(\xi) \times_{\Xi} \mathbf{A}(\xi))$  and  $\psi_{\xi\xi}(\mathbf{A}(\xi) \times_{\Xi} \mathbf{A}(\xi))$  covers together by the transition functions  $\Omega_{\xi,\xi}$ .

More specifically, the equivalence relations in the category of elements of such a structure sheaf, represented by a Boolean system of measurement localizations, have to be taken into account according to the coequalizer of coproduct representation of the colimit. Equivalence relations of this form give rise to congruences in the structure sheaf of Boolean coefficients, which are expressed categorically as a colimit in the category of elements of such a structure sheaf. In this perspective the generalized manifold, which represents categorically a quantum observable object, is understood as a colimit in a sheaf of Boolean coefficients, that contains compatible families of modeling Boolean observables. It is important to underline the fact that the organization of Boolean coordinatizing objects in localization systems takes the form of interconnection of these modeling objects through the categorical construction of colimit, the latter being the means to comprehend an object of complex structure (Quantum Observable) from simpler objects (Boolean Observables).

## 6 Isomorphic Representations of Quantum Observables by Boolean Localization Systems

The ideas developed in the previous section, may be used to provide the basis for the representation of Quantum observables and their associated Quantum event algebras, in terms of Boolean covering systems, if we pay attention to the counit of the established adjunction, denoted by the vertical map in [Diagram 12].

The diagram suggests that the representation of a quantum observable  $\Xi$

in  $\mathcal{O}_Q$ , and, subsequently, of a quantum event algebra  $L$  in  $\mathcal{L}$ , in terms of a coordinatization system of measurement localizations, consisting of Boolean coefficients, is full and faithful, iff the counit of the established adjunction, restricted to that system, is an isomorphism, that is, structure-preserving, 1-1 and onto.

The physical significance of this representation lies on the fact that, that the whole information content contained in a quantum event algebra, is preserved by every covering Boolean system, qualified as a system of measurement localizations. The preservation property is established by the counit isomorphism. It is remarkable that the categorical notion of adjunction provides the appropriate formal tool for the formulation of invariant properties, giving rise to preservation principles of a physical character.

If we return to the intended representation we realize that the surjective property of the counit guarantees that the Boolean domain covers, being themselves objects in the category of elements,  $\mathbf{G}(\mathbf{R}(L), B)$ , cover entirely the quantum event algebra  $L$ , whereas its injective property, guarantees that any two covers are compatible in a system of measurement localizations. Moreover, since the counit is also a homomorphism, it preserves the algebraic structure.

In the physical state of affairs, each cover corresponds to a set of Boolean events actualized locally in a measurement situation. The equivalence classes of Boolean domain covers represent quantum events in  $L$ , through compatible coordinatizations by Boolean coefficients. Consequently, the structure of a quantum event algebra is being generated by the information that its structure preserving maps, encoded as Boolean covers in measurement localization systems, carry, as well as their compatibility relations.

## 7 Implications for Quantum Logic

The covering process leads naturally to a contextual description of quantum events (or quantum propositions) with respect to Boolean reference frames of measurement, and finally to a representation of them as equivalence classes of unsharp Boolean events. The latter term is justified by the fact that, in case,  $L$  signifies a truth-value structure, each cover can be interpreted as an unsharp Boolean algebra of events, corresponding to measurement of observable  $\Xi$ . More concretely, since covers are maps  $[\psi_B]_\Xi : \mathbf{A}(B_\Xi) \longrightarrow L$ , each Boolean event realized in the domain  $B_\Xi$ , besides its true or false truth value assignment in a measurement context, related to the outcome of an experiment that has taken place, is also assigned a truth value representing its relational information content for the comprehension of the coherence of



the whole quantum structure, measured by the degrees in the poset  $L$ , or equivalently by the degrees assigned to its poset structure of localization systems.

Between these two levels of truth value assignment, there exists an intermediate level, revealed by the instantiation of the Boolean power construction in the context of the Grothendieck fibration technique. This intermediate level, refers to a truth value assignment to propositions describing the possible behavior of a quantum system in a specified Boolean context of observation, without, having passed yet, an experimental test.

We may remind that the fibration induced by a presheaf of Boolean algebras  $\mathbf{P}$ , provides the category of elements of  $\mathbf{P}$ , denoted by  $\mathbf{G}(\mathbf{P}, \mathcal{B})$ . Its objects are all pairs  $(B, p)$ , and its morphisms  $(\dot{B}, \dot{p}) \longrightarrow (B, p)$  are those morphisms  $u : \dot{B} \longrightarrow B$  of  $\mathcal{B}$  for which  $pu = \dot{p}$ . Projection on the second coordinate of  $\mathbf{G}(\mathbf{P}, \mathcal{B})$ , defines a functor  $\mathbf{G}_{\mathbf{P}} : \mathbf{G}(\mathbf{P}, \mathcal{B}) \longrightarrow \mathcal{B}$ . If  $B$  is an object of  $\mathcal{B}$ , the inverse image under  $\mathbf{G}_{\mathbf{P}}$  of  $B$  is simply the set  $\mathbf{P}(B)$ .

As we have explained, the objects of the category of elements  $\mathbf{G}(\mathbf{R}(L), B)$  constitute Boolean domain covers for measurement and have been identified as Boolean reference frames on a quantum observable structure.

We notice that the set of objects of  $\mathbf{G}(\mathbf{R}(L), \mathcal{B})$  consists of all the elements of all the sets  $\mathbf{R}(L)(B)$ , and more concretely, has been constructed from the disjoint union of all the sets of the above form, by labeling the elements. The elements of this disjoint union are represented as pairs  $(B, \psi_B : \mathbf{A}(B) \longrightarrow L)$  for all objects  $B$  of  $\mathcal{B}$  and elements  $\psi_B \in \mathbf{R}(L)(B)$ .

Taking into account the projection functor, defined above, this set is actually a fibered structure. Each fiber is a set defined over a Boolean algebra relative to which a measurement result is being coordinatized. If we denote by  $(\psi_B, q)$  the elements of each fiber, with  $\psi_B \in \mathbf{R}(L)(B)$  and  $q \in \mathbf{A}(B)$ , then the set of maps

$$(\psi_B, q) \longrightarrow q$$

can be interpreted as the Boolean power of the set

$$\Upsilon_B = \{(\psi_B, q), \psi_B \in \mathbf{R}(L)(B), q \in \mathbf{A}(B)\}$$

with respect to the underlying Boolean algebra  $B$  [14].

The Boolean power construction forces an interpretation of the Boolean algebra relative to which a measurement result is being coordinatized, as a domain of local truth values, with respect to, a measurement that has not taken place yet. Moreover the set of local measurement covers defined over  $B$ , is considered as a Boolean-valued set. In this sense, the local coordinates corresponding to a Boolean domain of measurement, may be considered as Boolean truth values.

We further observe that the set of objects of  $\mathbf{G}(\mathbf{R}(L), B)$  consists of the disjoint union of all the fibers  $\Upsilon_B$ , denoted by  $\Upsilon = \coprod_B \Upsilon_B$ . This set can also acquire a Boolean power interpretation as follows:

We define a binary relation on the set  $\Upsilon$  according to:

$$(\psi_{\dot{B}}, \hat{q}) \otimes (\psi_B, q) \text{ iff } \exists \eta : \psi_{\dot{B}} \longrightarrow \psi_B : \eta(\hat{q}) = q, \psi_{\dot{B}} = \psi_B \circ \eta.$$

It is evident that for any  $\eta : \dot{B} \longrightarrow B$  we obtain:  $(\psi_B \circ \eta, \hat{q}) \otimes (\psi_B, \eta(\hat{q}))$ . Furthermore, we require the satisfaction of the compatibility relations that are valid in a system of localizations. Then it is possible to define the Boolean power of the set  $\Upsilon$  with respect to the maximal Boolean algebra belonging to such a compatible system of localizations. We may say that the Boolean coordinates, interpreted as local Boolean truth values, via the Boolean power construction, reflect a relation of indistinguishability due to overlapping of the corresponding covers.

The viewpoint of Boolean valued sets, has far reaching consequences regarding the interpretation of quantum logic, and will be discussed in detail, in a future work. At the present stage, we may say that the logical interpretation of the Boolean fibration method, seems to substantiate Takeuti's and Davis's approach to the foundations of quantum logic [15, 16], according to whom, quantization of a proposition of classical physics is equivalent to interpreting it in a Boolean extension of a set theoretical universe, where  $B$  is a complete Boolean algebra of projection operators on a Hilbert space. In the perspective of the present analysis, we may argue that the fibration technique in the presheaf of Boolean algebras  $\mathbf{G}(\mathbf{R}(L), B)$ , provides the basis for a natural interpretation of the logic of quantum propositions, referring to the possible behavior of a quantum system in a concrete localization context with respect to an experimental test that has not been actualized yet, in terms of a truth value assignment, assuming existence in the corresponding Boolean context of a covering system, and realized in terms of local valuations on the Boolean coordinates of the specified cover.

## 8 Differential Geometry in the Quantum Regime

The Boolean manifold representation proposition permits the characterization of each quantum observable  $\Xi$  in  $\mathcal{O}_Q$  as a system of compatible measurement localizations consisting of Boolean observables, provided that the counit of the fundamental adjunction restricted to this system is an isomorphism [Diagram 12].

An important insight is provided by the application of the Stone representation theorem for Boolean algebras, according to which, it is legitimate to replace Boolean algebras by fields of subsets of a measurement space,

providing naturally an operationalization of the meaning of Boolean covers. If we replace each Boolean algebra  $B$  in  $\mathcal{B}$  by its set-theoretical representation  $[\Sigma, B_\Sigma]$ , consisting of a local measurement space  $\Sigma$  and its local field of subsets  $B_\Sigma$ , it is possible to define local measurement space charts  $(B_\Sigma, \psi_{B_\Sigma} : \mathbf{A}(B_\Sigma) \longrightarrow L)$  and corresponding space localization systems for quantum observable  $\Xi$  over quantum event algebra  $L$  in  $\mathcal{L}$ . Again from local measurement space charts  $(B_\Sigma, \psi_{B_\Sigma} : \mathbf{A}(B_\Sigma) \longrightarrow L)$  we may form their equivalence classes, which modulo the conditions for compatibility on overlaps, will represent a single quantum event in  $L$ . Under these circumstances, we may interpret the equivalence classes of local space charts  $\psi_{B_\Sigma} \otimes a$ ,  $a \in \mathbf{A}(B_\Sigma)$  as the experimental actualizations of the quantum events in  $L$ , corresponding to measurement of observables  $\Xi$ . The local measurement space charts  $(B_\Sigma, \psi_{B_\Sigma} : \mathbf{A}(B_\Sigma) \longrightarrow L)$  and  $(C_\Sigma, \psi_{C_\Sigma} : \mathbf{A}(C_\Sigma) \longrightarrow L)$  are compatible in a system of measurement localizations for observable  $\Xi$  defined over  $L$ , iff for some  $(D_\Sigma, \psi_{D_\Sigma} : \mathbf{A}(D_\Sigma) \longrightarrow L)$  in the system of localizations, and  $a \in \mathbf{A}(B_\Sigma)$ ,  $b \in \mathbf{A}(C_\Sigma)$ ,  $c, d \in \mathbf{A}(D_\Sigma)$ , the following conditions hold:

$$\psi_{B_\Sigma} \otimes a = \psi_{D_\Sigma} \otimes c$$

$$\psi_{C_\Sigma} \otimes b = \psi_{D_\Sigma} \otimes d$$

We note that we could equivalently consider the local space as a compact Hausdorff space, the compact open subsets of which are the maximal filters or the prime ideals of the underlying Boolean algebra.

The pullback compatibility condition, which is in injective correspondence with the one in  $\mathcal{L}$ , since, it holds in a localization system, may be interpreted in the operational framework, as denoting that, two local space representations of a quantum observable satisfy the compatibility condition on overlapping regions, iff their associated measurements are equivalent to measurements sharing the same experimental arrangement. We also observe that the inverse of a local space representation of a quantum observable plays the role of a random variable on this local space  $\Sigma$ . Consequently, every quantum observable may be considered locally, as a measurable function defined over the local measurement space  $\Sigma$ .

Phrased differently, random variables defined over local spaces provide Boolean coordinatizations for a quantum observable, and moreover, satisfy compatibility conditions on the overlaps of their local domains of definition. Subsequently, if we consider the category of these local spaces, the collection of measurable functions defined locally over them, provide a sheaf of Boolean coefficients, defined over the category of local spaces for the measurement of a quantum observable. The simplest case, is when, local spaces are partially ordered by inclusion. In this case, the sheaf of Boolean observable coordinates

can be considered as a sheaf of measurable functions defined locally over a topological space  $\Theta$ , being the union of the partially ordered local spaces. In the perspective of the interpretation put forward, the essence of a quantum observable is captured by a colimit in the category of elements of the sheaf of measurable functions, over the category of local spaces. The colimit in the category of elements of that sheaf, is expressed, as an equivalence relation on the collection of the locally defined Boolean coordinatizations according to the relation

$$pv \otimes \acute{q} = p \otimes v\acute{q}, \quad p \in \mathbf{R}(L)(\xi), \acute{q} \in \mathbf{A}(\acute{\xi}), v : \acute{\xi} \longrightarrow \xi$$

Consequently, a quantum observable is apprehended by an equivalence relation on the collection of its locally defined Boolean coordinatizations. Adopting the appropriate conditions for the existence of the topological space  $\Theta$  as above, a quantum observable is represented by a quotient construction, consisting of the sheaf of measurable functions defined locally over  $\Theta$ , modulo the ideal generated by the equivalence relations. This quotient construction on Boolean coordinatizations is a sheaf of equivalence classes of measurable functions on  $\Theta$ . Addition and multiplication over  $\mathbf{R}$  induce the structure of a sheaf of  $\mathbf{R}$ -algebras (or a sheaf of rings).

Instead of the quotient sheaf of measurable functions defined locally over  $\Theta$ , we could also consider the quotient sheaf of  $\mathbf{R}$ -algebras of continuous or smooth functions corresponding to local coordinatizations of a quantum observable. A natural question that arises in this setting, is if it could be possible, to consider the above quotient sheaf of  $\mathbf{R}$ -algebras as the structure algebra sheaf of a generalized space, corresponding to the category of local spaces, the simplest case being a structure algebra sheaf of the topological space  $\Theta$ . From a physical point of view, this move would reflect the appropriate generalization of the arithmetics, or sheaves of coefficients, that have to be used in the transition from the classical to the quantum regime. In the classical case, each local space endowed with appropriate topological and differential properties, may acquire the structure of a differentiable manifold, which, in turn, is characterized completely, by the structure sheaf of smooth functions, playing the equivalent role of a Boolean observer arithmetics. Since, all classical theoretical observables are always compatible, the sheaf of coefficients (smooth functions) can be used globally, giving rise to the differential geometric mechanism of smooth manifolds.

On the other side, we have seen that a quantum observable cannot be apprehended by the use of a single Boolean observer arithmetic, but, there is a necessity of employing a whole system of local arithmetics over a category of local spaces, which are constrained to specify appropriate compatibility

and equivalence relations, according to the fundamental adjunction of the categorical scheme. A single Boolean observer's arithmetic inevitably suppresses information about a quantum system and reflects a fuzzy apparatus for probing the structure of the quantum regime.

This observation is particularly enlightening when considering the notion of space, or even space-time, at the quantum regime. It naturally points to an understanding of the, so called, space-time manifold singularities, as reflections of the inability of a Boolean observer's arithmetic, consisting of smooth functions, to probe the quantum level of structure of this entity. According to the perspective of our discussion, the appropriate generalized arithmetic that would correspond, as a structure sheaf, to a quantum conception of space-time, would be the quotient sheaf of  $\mathbf{R}$ -algebras. This sheaf would contain compatible overlapping systems of local arithmetics, consisting of locally defined smooth functions over the category of local spaces, modulo the ideal corresponding to their equivalence relations, as a reflection of the interconnecting machinery of the colimit construction.

The appropriate framework to accommodate structure sheaves of the above form is Abstract Differential Geometry (ADG), developed by Mallios in [17], and discussed in relation to space-time singularities in [18]. ADG is an extension of classical Differential Geometry which does no longer use any notion of calculus. Instead of smooth functions, one starts with a sheaf of algebras, to be interpreted as a structure sheaf, considered on an arbitrary topological space, which serves as the base space of the sheaves involved. The important thing is that these sheaves of algebras, which in the perspective of the present categorical scheme, correspond to quantum observables, can be interrelated with appropriate differentials, instantiated as suitable Leibniz sheaf morphisms, and, constituting appropriate differential complexes. This interpretation is suited to the development of Differential Geometry in the regime of Quantum systems and will be carried out at a later stage. Most significantly, it emphasizes the thesis that the intrinsic mechanism of Differential Geometry is of an operational character, referring directly to the objects of enquiry, being in the proposed scheme, the quantum observables.

## 9 Conclusions

The conceptual roots of the proposed relativistic perspective on quantum structure, established by systems of Boolean measurement localization systems, is located on the physical meaning of the adjunction between presheaves of Boolean observables and quantum observables.

Let us consider that  $\mathbf{Sets}^{\mathcal{B}^{op}}$  is the universe of Boolean observable event

structures modeled in **Sets**, or else the world of Boolean windows, and  $\mathcal{L}$  that of Quantum event structures. In the proposed interpretation the functor  $\mathbf{L} : \mathbf{Sets}^{\mathcal{B}^{op}} \longrightarrow \mathcal{L}$  can be comprehended as a translational code from Boolean windows to the Quantum species of event structure, whereas the functor  $\mathbf{R} : \mathcal{L} \longrightarrow \mathbf{Sets}^{\mathcal{B}^{op}}$  as a translational code in the inverse direction. In general, the content of the information is not possible to remain completely invariant translating from one language to another and back. However, there remain two ways for a Boolean-event algebra variable set  $\mathbf{P}$ , or else Boolean window, to communicate a message to a quantum event algebra  $L$ . Either the information is given in Quantum terms with  $\mathbf{P}$  translating, which we can be represented as the quantum homomorphism  $\mathbf{LP} \longrightarrow L$ , or the information is given in Boolean terms with  $L$  translating, that, in turn, can be represented as the natural transformation  $\mathbf{P} \longrightarrow \mathbf{R}(L)$ . In the first case, from the perspective of  $L$  information is being received in quantum terms, while in the second, from the perspective of  $\mathbf{P}$  information is being sent in Boolean terms. The natural bijection then corresponds to the assertion that these two distinct ways of communicating are equivalent. Thus, the physical meaning of the adjoint situation signifies a two-way dependency of the involved languages in communication, with respect to the variation of the information collected in localization contexts of measurement. The adjunctive correspondence is the categorical formal tool needed at the level of relating relations. It specifies the necessary and sufficient conditions for establishing a notion of amphidromous dependent variation, realized operationally in the process of extraction of the information content enfolded in a quantum observable structure, through the pattern recognition characteristics of specified Boolean domain covers. More remarkably, the representation of a quantum observable as a categorical colimit, resulting from the same adjunctive relation, reveals an entity that can admit a multitude of instantiations, specified mathematically by different coordinatizing Boolean coefficients in Boolean localization systems.

The underlying invariance property specified by the adjunction is associated with the informational content of all these, phenomenally different instantiations in distinct measurement contexts, and can be formulated as follows: the informational content of a quantum observable structure remains invariant, with respect to Boolean domain coordinatizations, if and only if, the counit of the adjunction, restricted to covering systems, qualified as Boolean localization systems, is an isomorphism. Thus, the counit isomorphism, provides a categorical equivalence, signifying an invariance in the translational code of communication between Boolean windows, acting as localization devices for measurement, and quantum systems. Finally, the semantical aspects of that categorical equivalence, point towards investiga-

tions, regarding:

On the one side, the study of consequences of a naturally forced sheaf theoretical formulation of the logic of quantum propositions, and on the other, the development of an algebraic differential geometric machinery suited to the quantum level of observable structure, as has been discussed in the last two sections. It seems that the parallel development of the above research directions, on the basis of the physical meaning of the existing categorical equivalence, will substantiate a paraphrase of Lawvere's dictum [18], according to which:

Algebraic Quantum Geometry = Geometric Quantum Logic

**Acknowledgments:** I would gratefully like to acknowledge A. Mallios and S. Ivanov for many stimulating discussions. I would also wish to thank J. L. Bell, J.P. Marquis, V. Tsanov and A. Arrindell for their suggestions. The author is member of the EDGE Research Training Network HPRN-CT-2000-00101, supported by the European Human Potential Programme.

## References

- 1 F. W. Lawvere and S. H. Schanuel, *Conceptual Mathematics*, Cambridge University Press, Cambridge, 1997.
- 2 S. MacLane, *Categories for the Working Mathematician*, Springer-Verlag, New York, 1971.
- 3 S. MacLane and I. Moerdijk, *Sheaves in Geometry and Logic*, Springer-Verlag, New York, 1992.
- 4 J. L. Bell, *Toposes and Local Set Theories*, Oxford University Press, Oxford, 1988.
- 5 V. S. Varadarajan, *Geometry of Quantum Mechanics*, Vol. 1, Van Nostrand, Princeton, New Jersey, 1968.
- 6 G. Birkhoff and J. von Neumann, The logic of Quantum Mechanics, *Annals of Mathematics* **37**, 823 (1936).
- 7 S. Kochen and E. Specker, The problem of hidden variables in Quantum Mechanics, *J. Math. and Mech.* **17**, 59 (1967).
- 8 J. Butterfield and C. J. Isham, A Topos Perspective on the Kochen-Specker Theorem: I. Quantum States as Generalized Valuations, *International Journal of Theoretical Physics* **37**, 2669 (1998).
- 9 J. Butterfield and C. J. Isham, A Topos Perspective on the Kochen-Specker Theorem: II. Conceptual Aspects and Classical Analogues, *International Journal of Theoretical Physics* **38**, 827 (1999).
- 10 J. P. Rawling and S. A. Selesnick, Orthologic and Quantum Logic. Models and Computational Elements, *Journal of the Association for Computing Machinery* **47**, 721 (2000).



- 11 I. Raptis, Presheaves, Sheaves, and their Topoi in Quantum Gravity and Quantum Logic, gr-qc/0110064.
- 12 J. Butterfield and C. J. Isham, Some Possible Roles for Topos Theory in Quantum Theory and Quantum Gravity, *Foundations of Physics* **30**, 1707 (2000).
- 13 M. Artin, A. Grothendieck, and J. L. Verdier, *Theorie de topos et cohomologie etale des schemas*, (Springer LNM 269 and 270, Springer-Verlag, Berlin, 1972).
- 14 J. L. Bell, *Boolean-Valued Models and Independence Proofs in Set Theory*, Oxford University Press, Oxford, 1985.
- 15 G. Takeuti, Two Applications of Logic to Mathematics, *Mathematical Society of Japan* **13**, Kano Memorial Lectures 3 (1978).
- 16 M. Davis, A Relativity Principle in Quantum Mechanics, *International Journal of Theoretical Physics* **16**, 867 (1977).
- 17 A. Mallios, *Geometry of Vector Sheaves: An axiomatic Approach to Differential Geometry*, vols 1-2, Kluwer Academic Publishers, Dordrecht (1998).
- 18 A. Mallios, Remarks on Singularities, gr-qc/0202028.
- 19 F. W. Lawvere, Continuously Variable Sets: Algebraic Geometry=Geometric Logic, *Proceedings of the Logic Colloquium in Bristol*, North-Holland, Amsterdam (1975)